



Simulation tool for policymakers: evaluation of the socioeconomic effects of GHG reduction measures

Aleksejs Nipers, Kristīne Valujeva
Latvia University of Life Sciences and Technologies

LIFE OrgBalt, LIFE18 CCM/LV/001158

EU LIFE Programme project

“Demonstration of climate change mitigation potential of nutrients rich organic soils in Baltic States and Finland”



Latvia University of Life Sciences and Technologies



NATURAL RESOURCES INSTITUTE FINLAND



LITHUANIAN RESEARCH CENTRE FOR AGRICULTURE AND FORESTRY



Succow Stiftung






FOREST



ARABLE



ABANDONED



GRASSLAND



SCENARIO ANALYSIS

IMPACT

?

IMPACT


?

IMPACT

?

SIMULATION TOOL

- Simulation tool is data-based tool for policy planning and decision making at regional and national level.
- It evaluates the impact of climate change mitigation measures on socio-economic indicators and GHG emission reduction at national level for three Baltic States.
- Results of Simulation tool also shows possible spatial location of the GHG emission reduction measures.



Step 1: to collect agricultural and forestry data for the creation of a detailed land use data layer for each polygon.

Step 2: to collect data for organic soil.

Step 3: to generate working files for three Baltic States including only those agricultural and forestry areas located on organic soil

Step 4: to cut area from generated working files for each scenario based on predefined area selection criteria.


Step 5: impact assessment on profit, employment, and GHG emissions after implementation of scenarios.

SPATIAL DATA

SOIL TYPE



LAND USE





Organic soil (Yes/No)

ORTHOPHOTO




Agricultural land (crop, crop group, area, support, organic)
Forest land (forest type, main tree species, area, stand age, restrictions)
Protected areas (type, area, restrictions)






LVC301: CONVERSION OF CROPLAND TO GRASSLAND




Description	Area selection criteria	Land use after implementation
Cropland with nutrient-rich organic soil conversion to grassland. Increased carbon stock in soil and below-ground biomass, reduced risks of nutrient leaching and soil erosion.	Organic soil, arable land without perennial plantations	Grassland




LVC302: CONVENTIONAL AFFORESTATION (SPRUCE)

2030




2050




Description	Area selection criteria	Land use after implementation
Demonstration of the reduction of GHG emissions from area previously used as pasture or perennial grassland for fodder production by afforestation with spruce. Reduced GHG emissions from soil. Accumulation of CO ₂ in living and dead biomass, soil and litter and replacement effect of forest biofuel and harvested wood products. Shorter rotation and more intensified management ensures higher yield and replacement effect, as well as reduces carbon losses due to root rot and other disturbances.	Organic soil, grassland, perennial grassland, arable land without perennial plantations	Forest stand with spruce

Area of LVC302, ha




LVC303: INTRODUCTION OF FOREST PALUDICULTURE (DECIDIOUS TREES)

2030




2050




Description	Area selection criteria	Land use after implementation
Reduction of GHG emissions by establishing forest paludiculture (dominant species - black alder and birch) in grassland with nutrient-rich organic soil and increased groundwater level.	Organic soil, grassland, perennial grassland, arable land without perennial plantations	Forest stand with black alder and birch

Area of LVC303, ha



■ EE ■ LV ■ LT

LVC306: AGROFORESTRY – FAST GROWING TREES AND GRASS




Description	Area selection criteria	Land use after implementation
GHG emissions reduction through transformation of cropland to tree plantation. Projected reduction of GHG emissions is related to the decrease of N ₂ O and CO ₂ emissions from soil as well as to the increase of CO ₂ removals in living biomass and other carbon pools.	Organic soil, arable land without perennial grassland and perennial plantations	Forest stand with poplar

Area of LVC306, ha


0 50 000 100 000 150 000 200 000 250 000 300 000

■ EE ■ LV ■ LT

LVC310: FAST GROWING SPECIES IN RIPARIAN BUFFER ZONES




Description	Area selection criteria	Land use after implementation
GHG emissions reduction through transformation of strip areas along drainage ditches in cropland to tree plantation areas that avoid nutrient leaching and increase carbon removals in living biomass and other carbon pools. Projected reduction of GHG emissions is related to the decrease of N ₂ O and CO ₂ emissions from soil as well as to the increase of CO ₂ removals in living biomass and other carbon pools.	Organic soil, agricultural land, buffer zone at least 9.5 m wide from the edge of the ditch	Forest plantation with poplar and willow




LVC307: APPLICATION OF WOOD ASH IN SPRUCE TREE STANDS

2030



2050



Description

GHG emissions reduction in spruce stands on organic soils and lowered ground water table by implementation of wood ash after thinning thus enhancing stand growing conditions. Projected reduction of GHG emissions is related to groundwater level reduction, related to increase in growing stock increment and increased water amount used for transpiration processes – thus decreasing CH₄ emissions and increasing CO₂ removals in living biomass.


Area selection criteria

Organic soil, forest stand classification Kv, Km, Ks, Kp, II-IV site index, spruce at least 50%, age at least 20 years


Land use after implementation

Forest stand with spruce


Area of LVC307, ha




LVC308: CONTINUOUS FOREST IN SPRUCE STAND




Description	Area selection criteria	Land use after implementation
<p>GHG emissions reduction in spruce stand by replacing clear felling with selective felling. Projected reduction of GHG emissions is related to the increase of groundwater level in an alternative – clear felling scenario. Increase of groundwater level is associated with significant increase of CH₄. In the case of selective felling increase of groundwater levels should be smaller thus also increase of GHG emissions is smaller.</p>	<p>Organic soil, forest stand classification Pv, Nd, Db, Lk, Kv, Km, Ks, Kp, main specie spruce, age 81 years</p>	<p>Forest stand with spruce</p>




LVC309: FOREST REGENERATION WITH BLACK ALDER AND BIRCH IN NON-DRAINED ORGANIC SOIL




Description	Area selection criteria	Land use after implementation
<p>GHG emissions reduction in black alder and birch stand by using genetically selected planting material and improving hydrological regime. Projected reduction of GHG emissions is related to groundwater level stabilizing during forest regeneration phase and better growth conditions and increased CO₂ removals in forest biomass and other carbon stocks.</p>	<p>Organic soil, forest stand classification Pv, Nd, Db, Lk, main specie black alder, birch, age 71 years, I-III site index</p>	<p>Forest stand with black alder and birch</p>




LVC311: RIPARIAN BUFFER ZONE IN FOREST LAND PLANTED WITH BLACK ALDER




Description	Area selection criteria	Land use after implementation
<p>GHG emissions reduction in deciduous tree stands on organic soils with increased ground water table by enhancing tree growing conditions, using high quality planting material and preparing soil with mounding method including establishing of deep furrows for excess surface water drainage in spring time and after rainfalls. Projected reduction of GHG emissions is related to groundwater level reduction, related to establishment of deep furrows - as a result decreasing CH₄ emissions and increasing CO₂ removals in living biomass.</p>	<p>Organic soil, forest stand classification Ks, Kp, buffer zones of reclamation systems in forest lands</p>	<p>Forest stand with black alder</p>
<p>Area of LVC311, ha</p>  <p>0 10 000 20 000</p> <p>■ EE ■ LV ■ LT</p>		


LVC312: FOREST REGENERATION WITH PINE IN NON-DRAINED ORGANIC SOIL




Description	Area selection criteria	Land use after implementation
GHG emissions reduction in coniferous stands on organic soils and increased ground water table by application of forest regeneration with high quality coniferous planting material and by using mounding method for soil preparation. Projected reduction of GHG emissions is related to groundwater level reduction, related to establishment of deep furrows - as a result decreasing CH ₄ emissions and increasing CO ₂ removals in living biomass because of enhanced forest growing conditions.	Organic soil, forest stand classification Pv, Nd, Db, main species birch (age 71, II-V site index), aspen (age 41, site index II-V), black alder (age 71, II-V site index),	Forest stand with pine





LVC313: STRIP HARVESTING IN PINE STAND




Description	Area selection criteria	Land use after implementation
GHG emissions reduction in pine stand by replacing clear felling with strip harvesting. Projected reduction of GHG emissions is related to the increase of groundwater level in an alternative – clear felling scenario. Increase of groundwater level is associated with significant increase of CH ₄ . In the case of strip harvesting increase of groundwater levels should be smaller thus also increase of GHG emissions is smaller.	Organic soil, forest stand classification Kv, Km, Ks, Kp, main specie pine, age 101 years, I-III site index	Forest stand with pine





RESTORE1: GROWING BLUEBERRIES IN WETLANDS



Description	Area selection criteria	Land use after implementation
Conversion of former peat extraction sites to agricultural land where tall highbush blueberry <i>Vaccinium corymbosum</i> , or lowbush blueberry <i>Vaccinium angustifolium</i> are grown.	Former peat extraction field	Perennial plantation




RESTORE2: GROWING CRANBERRIES IN WETLANDS




Description	Area selection criteria	Land use after implementation
Conversion of former peat extraction sites to agricultural land where large cranberry <i>Vaccinium macrocarpon</i> is grown.	Former peat extraction field	Perennial plantation


Area of Restore2, ha




IMPACT ON GHG EMISSIONS IN 2050



IMPACT ON PROFITS IN 2050




IMPACT ON EMPLOYMENT IN 2050



APPLICATION OF SIMULATION TOOL

Scenarios Maps



Select scenario:

LVC301 - CONVERSION OF CROPLAND TO GRASSLAND

Description

Conversion of cropland with nutrient-rich organic soil to grassland. Increased carbon stock in soil and below-ground biomass, reduced risks of nutrient leaching and soil erosion.

Select year:

2030 2050

Estonia

Scenario area: 15 619 ha

GHG emissions: -17 181 t CO₂ eq.

Profit: -842 601 EUR

Employment: -41 400 hours

Latvia

Scenario area: 30 767 ha

GHG emissions: -33 844 t CO₂ eq.

Profit: -1 272 646 EUR

Employment: -145 800 hours

Lithuania

Scenario area: 40 764 ha

GHG emissions: -44 840 t CO₂ eq.

Profit: -1 727 904 EUR

Employment: -419 400 hours



Thank you!



www.orgbalt.eu



@orgbalt



@orgbalt



LIFE OrgBalt



orgbalt



orgbalt

The project "Demonstration of climate change mitigation potential of nutrients rich organic soils in Baltic States and Finland" (LIFE OrgBalt, LIFE18 CCM/LV/001158) has received funding from the LIFE Programme of the European Union and the State Regional Development Agency of Latvia.  www.orgbalt.eu

The information reflects only the LIFE OrgBalt project beneficiaries' view and the European Commission's Executive Agency for Small and Medium-sized Enterprises is not responsible for any use that may be made of the information contained therein.



Latvia University
of Life Sciences
and Technologies

